

# Trigeminal somatosensory evoked potentials in trichloroethylene-exposed workers

Chokri Mhiri, MD, Fakhher Choyakh, MD, Mohamed Ben Hmida, MD,  
Imed Feki, MD, Mounir Ben Messaoud, MD, Nouri Zouari, MD.

---

## ABSTRACT

**Objectives:** To assess the value of trigeminal somatosensory evoked potentials (TSEP) in evaluating toxicity of trichloroethylene to the trigeminal nerve.

**Methods:** Trichloroethylene (TCE), an excellent fat solvent largely used in industry, has a neurotoxic effect particularly on the trigeminal nerve. In March 2002, several TCE-exposed workers presented to Habib Bourguiba University Hospital, Sfax, Tunisia, for burned or numbness of the face. Twenty-three workers manipulating TCE underwent clinical, biological and toxicological assessment. Neurophysiologic study consisted of the measurement of the latencies and the amplitudes of TSEP in 23 exposed workers and 23 controls.

**Results:** Abnormal TSEP were observed in 6 workers with clinical evidence of trigeminal involvement and in 9 asymptomatic workers. We observed a significant positive correlation between duration of exposure and the N2 latency ( $r=0.5$ ,  $p<0.01$ ) and P2 latency ( $r=0.6$ ,  $p<0.02$ ). The sensitivity of TSEP was 65% and the specificity was 100%.

**Conclusion:** Trigeminal somatosensory evoked potentials are a useful tool in the diagnosis and management of chronic neurological complications of TCE intoxication. Since alterations of TSEP precede clinical symptoms, trigeminal nerve involvement can be demonstrated at the infra-clinical stage.

Neurosciences 2004; Vol. 9 (2): 102-107

---

**T**richloroethylene (TCE) is a halogenated hydrocarbon widely used in industry as a degreasing solvent since the first world war. Trichloroethylene is a neurotoxic and can produce selective trigeminal nerve analgesia; hence, it was used as therapy of trigeminal neuralgia (tic douloureux).<sup>1,2</sup> Many cases of cranial neuropathies from industrial exposure to TCE (metal degreasing, dry cleaning and so forth) or intentional inhalation of TCE were reported.<sup>3-8</sup> However, only few electrophysiological reports studied TCE-induced trigeminal neuropathies and no clear conclusions were available.<sup>9-13</sup> The aims of this study were to report our experience with TCE and to evaluate the usefulness of the trigeminal somatosensory evoked

potentials (TSEP) in this intoxication. We investigate the effect of TCE on amplitude and latencies of TSEP. We establish the normal control values for our own laboratory. Then, we measured TSEP in persons with occupational TCE exposure.

**Methods.** In March 2002, several phosphate industry workers presented to our out patient hospital for burned or numbness of the face. These workers were exposed to TCE, they spent up to 6 hours daily within tanks, they have to clean the inside walls of these tanks to be painted. For this task, they used TCE in such confined atmosphere without any ventilation. Protective masks and

---

From the Department of Neurology (Mhiri, Feki) and Functional Investigations (Choyakh, Zouari), Habib Bourguiba University Hospital, and the Departments of Nephrology (Ben Hmida) and Occupational Medicine (Ben Messaoud), Hedi Chaker University Hospital, Sfax, Tunisia.

Received 21st April 2003. Accepted for publication in final form 20th September 2003.

Address correspondence and reprint request to: Prof. Chokri Mhiri, Department Head, Department of Neurology, Habib Bourguiba University Hospital, CP 3.029-Sfax, Tunisia. Tel. +216 (74) 450581. Fax. +216 (74) 243427. E-mail: chokri.mhiri@rns.tn

safety advice are provided but not adopted by the workers due to heat and transpiration. All workers of this phosphate industry factory who have an occupational exposure to TCE for at least 2 years were included in the study. Using this unique criterion, we collected in the Department of Neurology a group of 23 males with a mean age of  $36.7 \pm 9$  years (range 20-56 years). They usually manipulate TCE, and the mean duration of exposure to this organic solvent was  $12.4 \pm 8.3$  years (range: 2-27 years). The non-exposed group consisted of 23 healthy males working in the same factory but not exposed to TCE. Their mean age was  $36.6 \pm 9.8$  years (range 24-54 years). None of the control group had a history of solvent exposure. These latter subjects were comparable to TCE exposed group in term of age and gender and were free from potential environmental, morbid or iatrogenic neuropathy. Exclusion criteria for both groups were diabetes mellitus, trigeminal neuralgia, alcohol abuse, multiple sclerosis or significant drug or medication histories.

**Baseline clinical evaluation.** All subjects were asked for burned or numbness of the face, visual disturbances, restlessness, concentration difficulty and fatigue. We also looked for mood changes such as euphoria and irritability; they are considered if they were concomitant to intensive exposure and were remarkable by patient's family. Neurological examination paid particular attention to the assessment of the cranial nerves. Quality of life and degree of disability was evaluated by the Karnovsky et al<sup>14</sup> performance score.

**Laboratory screening.** Standard biological investigations included blood cell count, erythrocyte sedimentation rate, fasting blood glucose, blood urea nitrogen, plasma creatinine, plasma cholesterol, plasma triglycerides, plasma bilirubin and hepatic enzymes (transaminase and  $\gamma$ -glutamyl transferase).

Toxicologic investigations: Measurement of urinary metabolites of TCE (trichloroethanol, trichloroacetic acid) was performed 3 times for each worker and usually on Friday, which is the last day of the week. Only the highest value is considered. Repeated careful measurements of TCE air contents were made.<sup>15,16</sup> In air atmosphere, the maximum tolerated concentration of trichloroethylene was 50 ppm and in urine, the maximum tolerated concentration trichloroacetic acid 100 mg/g of creatinine.<sup>17,18</sup>

**Neurophysiological screening.** Trigeminal somatosensory evoked potentials were recorded using Nihon-Kohden EMG-evoked potential system (Tokyo, Japan) as described by Stöhr and Petrüch.<sup>19</sup> Every subject underwent consecutive bilateral recording of somatosensory evoked potentials following trigeminal stimulation. A surface electrode, applied 1 cm under the corner of the mouth, delivered square wave pulses of 0.1 ms

duration at the rate of 3/seconds. Stimulus intensity was adjusted to produce moderate twitching of the inferior lip. The recording electrode is placed over the contralateral parietal area using a non-cephalic reference. Evoked potentials were amplified and filtered between 20-2000 Hz. Averaging was performed on at least 500 responses over an analysis time of 100 ms. Two averages were superimposed to confirm reproductivity. Latency and amplitude were determined for each wave: first negative potential (N1), first positive potential (P1), second negative potential (N2), second positive potential (P2) and third negative potential (N3). Trigeminal somatosensory evoked potentials latencies higher than our control mean plus 2.5 SD are considered abnormal. Therefore, subjects with asymmetrical amplitudes of TSEP were considered as pathological (difference between the 2 sides was >50%). Motor nerve conduction velocities were measured for median, ulnar, medial and lateral popliteal nerves. For sensory nerve conduction velocities, lateral sural cutaneous nerve and median nerve were assessed.

**Statistical analyses.** Statistical methods included paired or unpaired Student's t-test as appropriate. Results are expressed as mean  $\pm$  1 standard deviation; *p* values <0.05 were considered significant. Spearman rank-correlation procedure was used for correlation analysis.

**Results. Clinical results (Table 1).** Among the 23 workers, only 6 presented sensory disturbance of the face. It consisted of numbness, burned, or both, of the cheeks and hypoesthesia in the territory of the first and second branch of the trigeminal nerve including corneal hypesthesia. Corneal and blink reflexes were absent in 2 (case 11-12) and weak in one (case 7). Moreover, general symptoms were observed usually during episodes of intensive exposure including fatigue (19 cases), consciousness disturbance (18 cases), stance and gait instability (15 cases) and disturbed behavior such as restlessness, irritability, concentration difficulty with euphoria (12 cases). These disturbances were transient, workers had a quiet normal activity and the Karnovsky performance score was high (80-100, mean of  $86 \pm 8$ ).

**Results of laboratory screening.** Blood cell count was normal. Fasting blood glucose was at  $4.8 \pm 0.2$  mmol/l (range 4-6), blood urea nitrogen at  $5 \pm 0.3$  mmol/l (range 4-7.2), plasma creatinine level  $123.8 \pm 12.7$   $\mu$ mol/l (range 70-251), plasma cholesterol  $5 \pm 0.3$  mmol/l (range 3.9-6.9), plasma triglyceride  $1.7 \pm 0.1$  mmol/l (range 1.1-2.7), plasma bilirubin  $10.2 \pm 1.1$   $\mu$ mol/l (range 6-17), alanine amino-transferase at  $24.1 \pm 2.8$  UI/l (range 13-45), asparagine amino-transferase at  $26.9 \pm 1.9$  UI/l (range 16-42) and  $\gamma$ -glutamyl transferase  $19.8 \pm 4.8$  UI/l (range 11-75). Mean urinary

trichloroethanol expressed as mg/g of urinary creatinine was  $79.3 \pm 8.8$ , trichloroacetic acid was  $32.6 \pm 4.6$  and total trichlorides was  $111.9 \pm 11.5$  (Table 2). Only one subject had urinary TCE metabolites levels over tolerated limits (case 2). Trichloroethylene air contents were over tolerated limits of 50 ppm and ranged from 50-150 ppm.

**Neurophysiological findings.** In the non-exposed group, N1 and P1 potentials were inconstant with low amplitude and frequently masked by the stimulation artifact. Whereas N2 potential was constant ample and was considered as the best component for analysis. On the other hand, P2 and N3 potentials were relatively constant but their latencies showed a large variability.

The latencies of N1 were  $12.7 \pm 1.1$  ms, P1 were  $18.2 \pm 1$  ms, N2 were  $27 \pm 1.1$  ms, P2 were  $35 \pm 2.7$  ms and N3 were  $45.9 \pm 3.8$  ms (Figure 1a). The amplitudes of the different components showed large variations in this group, indeed mean values of N1 were  $0.9 \pm 0.3$   $\mu$ v, P1 were  $1.1 \pm 0.4$   $\mu$ v, N2 were  $2.2 \pm 1.1$   $\mu$ v, P2 were  $1.1 \pm 0.4$   $\mu$ v and N3 were  $1.4 \pm 0.7$   $\mu$ v. No difference was observed between the 2 sides regarding latencies. Amplitude may vary from one side to other, but the difference did not exceed 50%. In the exposed group, mean

latencies of the different potentials were significantly delayed. However, a significant decrease of the amplitude was noted only for P1 and N2 potentials (Table 3). Delayed latencies were observed in 12 cases, they were unilateral in 9 cases and bilateral in 3 cases (Figure 1b). Asymmetrical amplitude was the only abnormality in 4 patients. The limb motor and sensory nerve conduction velocities were normal in all exposed workers. A significant positive correlation was found between duration of exposure and both N2 latency ( $r=0.5$ ,  $p<0.01$ ) and P2 latency ( $r=0.6$ ,  $p<0.02$ ) (Figure 2). No correlation was noted between the TSEP and urinary TCE metabolites.

**Discussion.** Industrial surveys have revealed a number of examples of isolated and otherwise, unexplained facial numbness in workers exposed to TCE. Selective damage of the fifth cranial nerve has been documented by autopsy.<sup>3</sup> The most striking changes were seen in the brainstem particularly the fifth cranial nerve nuclei, spinal tract and nerve roots. Paradoxically, the structures involved in TCE intoxication have no feature in common such as axon length and size, energy consumption rates,

**Table 1** - Clinical data of trichloroethylene-exposed workers.

Case	Age*	Duration of exposure (years)†	General symptoms				Trigeminal nerve involvement symptoms					
			Fatigue	Disturbed behavior	Consciousness disturbance	Gait instability	Karnovsky score‡	Numbness of the face	Hypesthesia of the face	Corneal hypesthesia	Corneal reflex	Blink reflex
1	30	9	+	+	+	+	80	-	-	-	+	+
2	20	3	-	-	+	+	90	-	-	-	+	+
3	27	5	+	+	+	+	80	-	-	-	+	+
4	47	30	+	+	+	-	80	+	-	-	+	+
5	34	5	+	-	+	-	90	-	-	-	+	+
6	34	15	+	+	+	+	80	+	+	+	+	+
7	42	17	+	+	+	-	80	+	+	+	+/-	+/-
8	51	25	+	-	+	+	90	-	+	+	+	+
9	48	16	+	+	+	+	80	+	+	+	+	+
10	35	12	+	+	+	+	80	-	-	-	+	+
11	46	27	+	+	+	+	80	+	+	+	-	-
12	41	21	+	+	+	+	80	+	+	+	-	-
13	31	13	-	-	-	-	100	-	-	-	+	+
14	56	22	+	-	+	-	90	-	-	-	+	+
15	29	4	+	+	+	+	80	-	-	-	+	+
16	37	6	+	-	+	+	80	-	-	-	+	+
17	28	6	+	-	+	+	80	-	-	-	+	+
18	33	13	+	+	+	+	80	-	-	-	+	+
19	40	5	+	+	+	+	80	-	-	-	+	+
20	29	6	+	-	-	-	100	-	-	-	+	+
21	30	2	+	-	-	+	90	-	-	-	+	+
22	26	7	-	-	-	-	100	-	-	-	+	+
23	39	17	-	-	-	-	100	-	-	-	+	+

\* mean  $\pm$  SD  $36.7 \pm 9$ , range 20 - 56 years, † mean  $\pm$  SD  $12.4 \pm 8$ , range 2 - 27 years, ‡ mean  $\pm$  SD  $86 \pm 8$ , range 80 - 100 years, + = presence, - = absence, +/- = present but weak


**Table 2** - Trichloroethylene urinary metabolites expressed as mg per gram of excreted creatinine.

Case	Trichloro-ethanol (mg/g of UC)	Trichloro-acetic acid (mg/g of UC)	Total trichlorides (mg/g of UC)	UC (g/l)
1	145	46	191	1.47
2	211	58	269	0.93
3	61	97	158	0.95
4	49	12	61	1.47
5	135	65	200	0.89
6	75	45	120	1.32
7	9.5	3	12.5	2.42
8	100	20	120	1.56
9	93	45	138	1.30
10	50	55	105	1.72
11	63	20	83	1.83
12	75	30	105	1.34
13	86	22	108	0.86
14	66	3	69	1.02
15	90	32	122	1.71
16	72	38	110	0.93
17	58	18	76	1.10
18	75	27	102	2.26
19	59	12	71	1.09
20	67	19	86	1.43
21	10	20	30	2.15
22	83	35	118	1.93
23	91	29	120	1.42
Mean $\pm$ SD	79.3 $\pm$ 42	32.6 $\pm$ 22	111.9 $\pm$ 55	1.4 $\pm$ 0.5
Range	9.5 - 211	3 - 97	12.5 - 269	0.9 - 2.4

UC - urinary creatinine

specific transmitter, metabolism and so forth. However, vascular permeability of the region accounts for the vulnerability of the trigeminal nerve nuclei and may explain specific damage.<sup>20</sup> On the other hand, the occurrence of orofacial herpes simplex lesions associated with TCE-induced trigeminal neuropathy suggests that TCE may act as a reactivating factor of the neurotropic virus.<sup>21</sup> The recording of TSEP following trigeminal stimulation is rarely used in current practice. Trigeminal somatosensory evoked potentials were first reported in humans by Stöhr and Petruch<sup>9</sup> and followed by Bennett and Jannetta,<sup>22</sup> Findler and Feinsod,<sup>23</sup> and by Garrel et al.<sup>24</sup> Clinical application was emphasized in trigeminal neuralgia<sup>25,26</sup> and TCE intoxication.<sup>9,10</sup> Stimulation can be applied on the lips<sup>10,19,22-24</sup> or on the tongue.<sup>26</sup>


In our outpatient hospital, several workers presented with trigeminal sensory disturbances. Anamnesis confirmed chronic TCE exposure. A cross-sectional clinical, toxicological and neurophysiological investigation was performed. In the studied group, only 6 of 23 had clinical evidence of trigeminal involvement. However, neurophysiological study disclosed abnormal TSEP in 15 subjects (65%). The abnormalities observed were similar to those reported by Garrel and Barret<sup>9</sup> and Dogui et al,<sup>10</sup> and it consists of delayed latencies, asymmetrical amplitude, or both. Unlike previous reports, we found a high percentage of abnormal TSEP in our exposed group (65%),



**Figure 1** - Graphs of trigeminal somatosensory evoked potentials in **a)** controls and **b)** trichloroethylene-exposed workers.

**Table 3** - Mean latencies and amplitudes of the trigeminal somatosensory evoked potentials in trichloroethylene (TCE) exposed workers and controls.

Potential	TCE-exposed	p value	Control
<b>N1</b>			
Latency (ms)	15.9 $\pm$ 1.7	<0.001	12.6 $\pm$ 0.9
Amplitude ( $\mu$ v)	1.0 $\pm$ 0.4	NS	1.0 $\pm$ 0.2
<b>P1</b>			
Latency (ms)	20.2 $\pm$ 2.5	<0.01	18.4 $\pm$ 1
Amplitude ( $\mu$ v)	0.8 $\pm$ 0.5	<0.02	1.1 $\pm$ 0.3
<b>N2</b>			
Latency (ms)	29.0 $\pm$ 3.5	<0.01	27.0 $\pm$ 0.9
Amplitude ( $\mu$ v)	1.6 $\pm$ 0.7	<0.05	2.3 $\pm$ 0.7
<b>P2</b>			
Latency (ms)	38.0 $\pm$ 4.6	0.01	35.0 $\pm$ 2.1
Amplitude ( $\mu$ v)	1.3 $\pm$ 0.6	NS	1.4 $\pm$ 1
<b>N3</b>			
Latency (ms)	52.4 $\pm$ 8.2	0.001	45.5 $\pm$ 3
Amplitude ( $\mu$ v)	1.7 $\pm$ 0.7	NS	1.4 $\pm$ 0.6
NS - not significant			



**Figure 2** - Positive correlation between duration of exposure and the N2 and P2 latencies.

whereas the percentage of positivity found by Garrel and Barret<sup>9</sup> was 38% and by Dogui et al<sup>10</sup> 45%. This difference may be explained by the long exposure duration in our series ( $12.4 \pm 8$  years). The most striking finding was a positive correlation between TSEP latencies and the exposure duration (for N2 latency  $r=0.5$  and  $p<0.01$  and for P2 latency  $r=0.6$  and  $p<0.02$ ), which may suggest a cumulative dose-response relationship. No correlation was found between neurophysiological alteration and urinary TCE metabolite concentration and TCE air-content. To assess the general validity of the TSEP measurement as an indicator of exposure to TCE, a 2 x 2 table was constructed (**Table 4**). The sensitivity was 65% and specificity was 100%.<sup>27</sup> The sensitivity of TSEP may be improved if the 3 branches of the trigeminal nerve are explored. Early evoked potentials can be recorded on the scalp after stimulation of the infra-orbital nerve. The waves have been named  $W_1$  with mean latency of 0.98 ms,  $W_2$  with mean latency of 1.92 ms and  $W_3$  with mean latency of 2.57 ms.<sup>28,29</sup> These early components are probably generated by the proximal part of the maxillary nerve or from the gasserian ganglion ( $W_1$ ) and from the trigeminal root fibres running through the brainstem before ( $W_2$ ) and after the dichotomisation ( $W_3$ ).<sup>30</sup> They are interesting to evaluate the trigeminal system in its peripheral parts, but it needs a very short pulse duration (0.05 ms) and alternating polarity essential for artifact suppression.<sup>31</sup> Middle latency components of TSEP (N1, P1, N2, P2 and N3) were considered to arise from supratthalamic generators. Since, the most neuropathological changes induced by TCE were seen in the brainstem especially the fifth nerve nuclei, it is preferable to study middle than early latency trigeminal evoked potentials. The blink reflex was reported as an alternative approach in the management of TCE exposed workers.<sup>11</sup> Although this technique seems to be easier than TSEP, it is less sensitive (50%) and specific (90%). The lack of sensitivity may be attributed to the fact that the test does not investigate trigeminal involvement specifically but facial and trigeminal nerve involvement.

**Table 4** - Matrice 2 by 2 of validity indicators regarding the trigeminal somatosensory evoked potentials (TSEP).

TSEP	Trichloroethylene exposed	Control
Normal TSEP	8	18
Pathologic TSEP	15	0

In conclusion, our study demonstrates that TSEP, a safe and non invasive investigation, is a useful tool in diagnosis and the management of trigeminal nerve involvement secondary to chronic TCE intoxication. Trigeminal somatosensory evoked potential abnormalities precede clinical symptoms and are related to the exposure duration. They should be performed periodically in all patients exposed to this organic solvent to detect neurological effects early.

**Acknowledgment.** We are deeply indebted to Dr. Romain K. Gherardi for critically reading and checking the English of this manuscript.

## References

- Defalque RJ. The specific analgesic effect of trichloroethylene upon the trigeminal nerve. *Anesthesiology* 1961; 22: 379-384.
- Seppäläinen AM. Halogenated hydrocarbons. In: Blum K, Manzo L, editors. Neurotoxicology. New York (NY): Marcel Dekker Inc; 1985. p. 459-472.
- Buxton PH, Hayward M. Polyneuritis cranialis associated with industrial trichloroethylene poisoning. *J Neurol Neurosurg Psychiatry* 1967; 30: 511-518.
- Mitchell ABS, Pearsons-Smith BG. Trichloroethylene neuropathy. *Br Med J* 1969; 1: 422-423.
- Lawrence WH, Partyka EK. Chronic dysphagia and trigeminal anesthesia after trichloroethylene exposure. *Ann Med Interna* 1981; 95: 710.
- Barret L, Faure J, Guiland B, Chomat D, Didier B, Debru JL. Trichloroethylene occupational exposure: elements for better prevention. *Int Arch Occup Environ Health* 1984; 53: 283-289.
- Szlatenyi CS, Wang RY. Encephalopathy and cranial nerve palsies caused by intentional trichloroethylene inhalation. *Am J Emerg Med* 1996; 14: 464-466.
- Kaneko T, Wang PY, Sato A. Assessment of the health effects of trichloroethylene. *Ind Health* 1997; 35: 301-324.
- Garrel S, Barret L. Potentiels évoqués somesthésiques du trijumeau dans les intoxications chroniques par le trichloréthylène. *Rev Electroencephalogr Neurophysiol Clin* 1986; 16: 133-140.
- Dogui MN, Mrizak M, Yacoubi B, Ben Hadj A, Paty J. Potentiels évoqués somesthésiques du trijumeau chez les travailleurs manipulant le trichloréthylène. *Neurophysiol Clin* 1991; 21: 95-103.
- Feldman RG, Niles C, Proctor SP, Jabre J. Blink reflex measurement of effects of trichloroethylene exposure on the trigeminal nerve. *Muscle Nerve* 1992; 15: 490-495.

12. Lash TL, Green LC. Blink reflex measurement of effects of trichloroethylene exposure on the trigeminal nerve. *Muscle Nerve* 1993; 16: 217-220.
13. Leandri M, Schizzi R, Scielzo C, Favale E. Electrophysiological evidence of trigeminal root damage after trichloroethylene exposure. *Muscle Nerve* 1995; 18: 467-468.
14. Karnofsky DA, Abelmann WH, Kraver LF, Burchenal JH. The use of nitrogen mustards in the palliative treatment of carcinoma. *Cancer* 1948; 634-656.
15. Jentsch D, Fraser AM. A laboratory evaluation of long term detector tubes: benzene, toluene, and trichloroethylene. *Am Ind Hyg Assoc J* 1981; 42: 810-823.
16. Boudène CL, Turbiaux M, Cluet JL, Truhaut R. Contribution à la fixation d'une concentration limite tolérable de trichloréthylène dans l'atmosphère des locaux de travail. *Arch Mal Prof* 1983; 44: 75-91.
17. Ertle T, Henschler D, Müller G, Spassowski M. Metabolism of trichloroethylene in man: the significance of trichloroethanol in long term exposure condition. *Arch Toxikol* 1972; 29: 171.
18. Vincent M, Faivre-Pierret R, Marka C, Vincent F, Benoît-Guyot JL, Faure J. La surveillance de l'exposition au trichloréthylène en milieu de travail. *Arch Mal Prof* 1980; 41: 189-194.
19. Stöhr M, Petrüch F. Somatosensory evoked potentials following stimulation of the trigeminal nerve in man. *J Neurol* 1979; 220: 95-98.
20. Jacobs JM. Vascular permeability and neurotoxicity. *Environ Health Perspect* 1978; 26: 107-116.
21. Cavanagh JB, Buxton PH. Trichloroethylene cranial neuropathy: is it really a toxic neuropathy or does it activate latent herpes virus. *J Neurol Neurosurg Psychiatry* 1989; 52: 297-303.
22. Bennett MH, Jannetta PJ. Trigeminal evoked potentials in humans. *Electroencephalogr Clin Neurophysiol* 1980; 48: 517-526.
23. Findler G, Feinsod M. Sensory evoked response to electrical stimulation of trigeminal nerve in humans. *J Neurosurg* 1982; 56: 545-549.
24. Garrel S, Barret L, Maynard R. Potentiel évoqués somesthésiques du trijumeau chez l'homme - premiers résultats. *Rev Electroencephalogr Neurophysiol Clin* 1982; 12: 129-134.
25. Stöhr M, Petrüch F, Scheglman K. Somatosensory evoked potentials following trigeminal nerve stimulation in trigeminal neuralgia. *Ann Neurol* 1981; 9: 63-66.
26. Altenmüller E, Cornelius CP, Buettner UW. Somatosensory evoked potentials following tongue stimulation in normal subjects and patients with lesions of the afferent trigeminal system. *Electroencephalogr Clin Neurophysiol* 1990; 77: 403-415.
27. Mausner JS, Bahn S. Epidemiology - An Introduction Text. 2nd ed. Philadelphia (PA): Saunders; 1985. p. 214-238.
28. Leandri M, Parodi CI, Favale E. Early evoked potentials detected from the scalp of man following infraorbital nerve stimulation. *Electroencephalogr Clin Neurophysiol* 1985; 62: 99-107.
29. Leandri M, Parodi CI, Favale E. Normative data on scalp responses evoked by infraorbital nerve stimulation. *Electroencephalogr Clin Neurophysiol* 1988; 71: 415-421.
30. Leandri M, Campbell JA. Origin of early waves evoked by infraorbital nerve stimulation in man. *Electroencephalogr Clin Neurophysiol* 1986; 65:13-19.
31. Soustiel JF, Feinsod M, Hafner H. Short latency trigeminal evoked potentials: normative data and clinical correlations. *Electroencephalogr Clin Neurophysiol* 1991; 80: 119-125.